Asthma Treatment Plan – Student

(This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) (Physician's Orders)

www.pacnj.org







(Please Print)		www.pai	og.org		
Name		Date of Birth	Effective Date		
Doctor	Parent/Guardian (if a	Parent/Guardian (if applicable)		Emergency Contact	
Phone	Phone	Phone		Phone	
HEALTHY (Green Zone)	Take daily control n	a "spacer" – use i	if directed.	Triggers Check all items that trigger	
You have all of these: Breathing is good No cough or wheeze Sleep through the night Can work, exercise, and play	MEDICINE HOW MUCH to take and HOW OFTEN to take it Advair® HFA		patient's asthma: Colds/flu Exercise Allergens Dust Mites, dust, stuffed animals, carpet Pollen - trees, grass, weeds Mold		
	None		fter taking inhaled medicineminutes before exercise.	SHIUKE	
You have any of these: Cough Mild wheeze Tight chest Coughing at night Other: If quick-relief medicine does not help within 15-20 minutes or has been used more than times and symptoms persist, call your doctor or go to the emergency room. And/or Peak flow from to	Continue daily control medicine(s) and ADD quick-relief medicine(s). MEDICINE HOW MUCH to take and HOW OFTEN to take it Combivent® Maxair® Xopenex® 2 puffs every 4 hours as needed Ventolin® Pro-Air® Proventil® 2 puffs every 4 hours as needed Albuterol 1.25, 2.5 mg 1 unit nebulized every 4 hours as needed Duoneb® 1 unit nebulized every 4 hours as needed Xopenex® (Levalbuterol) 0.31, 0.63, 1.25 mg 1 unit nebulized every 4 hours as needed Increase the dose of, or add: Other If quick-relief medicine is needed more than 2 times a week, except before exercise, then call your doctor.			products, scented products Smoke from burning wood, inside or outside Weather Sudden temperature change Extreme weather hot and cold Ozone alert days Foods:	
Your asthma is getting worse fast: • Quick-relief medicine did not help within 15-20 minut • Breathing is hard or fast • Nose opens wide • Ribs sho • Trouble walking and talking • Lips blue • Fingernails blue • Other: • Other:	Asthma can be a li MEDICINE Combivent® Maxair® Decemble Pro-Air® Pro-Air® Duoneb®	HOW MUCH to t Xopenex® oventil® g	ness. Do not wait! take and HOW OFTEN to take it 2 puffs every 20 minutes	Other: O This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs	
This is the control of the control o	sion to Self-administer Medication student is capable and has been instructed a proper method of self-administering of the nebulized inhaled medications named above cordance with NJ Law. student is not approved to self-medicate.	PARENT/GUARDIAN SIGNATI	URE		

Make a copy for parent and for physician file, send original to school nurse or child care provider.